



RESISTANCE TO DAMAGE, PESTS AND DISEASES	LOW	→							HIGH
Potato cyst nematode (<i>Globodera pallida</i> Pa 2/3,1)		2							
Potato cyst nematode (<i>Globodera rostochiensis</i> Ro1)		2							
Late blight on foliage (<i>Phytophthora infestans</i>)						5			
Late blight on tubers (<i>Phytophthora infestans</i>)		2							
Blackleg (<i>Pectobacterium atrosepticum</i>)								7	
Potato leafroll virus	1								
Potato virus Yo				4					
Skin spot (<i>Polyscytalum pustulans</i>)								8	
Common scab (<i>Streptomyces scabiei</i>)					5				
Powdery scab (<i>Spongospora subterranea</i>)					5				
Silver scurf (<i>Helminthosporium scabiei</i>)						6			
Black dot (<i>Colletotrichum coccodes</i>)				4					
Black scurf (<i>Rhizoctonia solani</i>)					5				
Dry rot (<i>Fusarium coeruleum</i>)				4					
Dry rot (<i>Fusarium sulphureum</i>)								9	
Internal damage (bruising)							7		
External damage (and splitting)							7		

FIELD AND STORE QUALITIES	LOW	→							HIGH
Drought resistance							7		
Heat tolerance						6			
Storability						6			