

| RESISTANCE TO DAMAGE, PESTS AND DISEASES | LOW | → | | | | | | | HIGH |
|--|------------------|---|---|---|---|---|---|---|------|
| Potato cyst nematode (<i>Globodera pallida</i> Pa 2/3,1) | | | 3 | | | | | | |
| Potato cyst nematode (<i>Globodera rostochiensis</i> Ro1) | | | | | | | | 8 | |
| Late blight on foliage (<i>Phytophthora infestans</i>) | | | | | 5 | | | | |
| Late blight on tubers (<i>Phytophthora infestans</i>) | | | 3 | | | | | | |
| Blackleg (<i>Pectobacterium atrosepticum</i>) | | | | | | 6 | | | |
| Potato leafroll virus | | 2 | | | | | | | |
| Potato virus Yo | | 2 | | | | | | | |
| Potato virus Ya | | | | | | | | | 9 |
| Potato virus Yn | 1 | | | | | | | | |
| Mop-top virus (Spraing) (PMTV) | | | | 4 | | | | | |
| Common scab (<i>Streptomyces scabiei</i>) | | | | 4 | | | | | |
| Powdery scab (<i>Spongospora subterranea</i>) | | | | | | | | 7 | |
| Silver scurf (<i>Helminthosporium scabiei</i>) | | | | | | | 6 | | |
| Black dot (<i>Colletotrichum coccodes</i>) | | | | | 5 | | | | |
| Black scurf (<i>Rhizoctonia solani</i>) | Awaiting results | | | | | | | | |
| Dry rot (<i>Fusarium coeruleum</i>) | | | | | | | | | 8 |
| Dry rot (<i>Fusarium sulphureum</i>) | | | | | | | | | 8 |
| Internal damage (bruising) | | | | | | | 6 | | |
| External damage (and splitting) | | | | | | | 6 | | |

| FIELD AND STORE QUALITIES | LOW | → | | | | | | | HIGH |
|---------------------------|-----|---|--|--|---|--|--|---|------|
| Drought resistance | | | | | 5 | | | | |
| Heat tolerance | | | | | 5 | | | | |
| Storability | | | | | | | | 7 | |