

# Caledonian Pearl

Early



RESISTANCE TO DAMAGE, PESTS AND DISEASES	LOW	→										HIGH
Potato cyst nematode ( <i>Globodera pallida</i> Pa 2/3,1)												2
Potato cyst nematode ( <i>Globodera rostochiensis</i> Ro1)	1											
Late blight on foliage ( <i>Phytophthora infestans</i> )												4
Late blight on tubers ( <i>Phytophthora infestans</i> )												5
Blackleg ( <i>Pectobacterium atrosepticum</i> )												6
Potato leafroll virus												6
Potato virus Yo	Awaiting results											
Potato virus Ya												9
Potato virus Yn												4
Skin spot ( <i>Polyscytalum pustulans</i> )	Awaiting results											
Common scab ( <i>Streptomyces scabiei</i> )	Awaiting results											
Powdery scab ( <i>Spongospora subterranea</i> )												8
Silver scurf ( <i>Helminthosporium scabiei</i> )	Awaiting results											
Black dot ( <i>Colletotrichum coccodes</i> )	Awaiting results											
Black scurf ( <i>Rhizoctonia solani</i> )	Awaiting results											
Dry rot ( <i>Fusarium coeruleum</i> )												4
Dry rot ( <i>Fusarium sulphureum</i> )												3
Internal damage (bruising)												7
External damage (and splitting)												6

FIELD AND STORE QUALITIES	LOW	→										HIGH
Drought resistance												7
Heat tolerance												7
Storability												7

**Disclaimer:** The contents are based on the best information available, but the selling parties will not accept responsibility for any inaccuracy caused by circumstances outside their control, nor will they be liable for any costs, loss, damage or injury resulting from interpretation of, or decisions based on, the information provided.