

Caledonian Rose

Maincrop



RESISTANCE TO DAMAGE, PESTS AND DISEASES	LOW	→								HIGH
Potato cyst nematode (<i>Globodera pallida</i> Pa 2/3,1)		2								
Potato cyst nematode (<i>Globodera rostochiensis</i> Ro1)										9
Late blight on foliage (<i>Phytophthora infestans</i>)					5					
Late blight on tubers (<i>Phytophthora infestans</i>)			3							
Blackleg (<i>Pectobacterium atrosepticum</i>)									7	
Potato leafroll virus										8
Potato virus Yo	Awaiting results									
Potato virus Ya										9
Potato virus Yn									7	
Skin spot (<i>Polyscytalum pustulans</i>)	Awaiting results									
Common scab (<i>Streptomyces scabiei</i>)									6	
Powdery scab (<i>Spongospora subterranea</i>)									6	
Silver scurf (<i>Helminthosporium scabiei</i>)	Awaiting results									
Black dot (<i>Colletotrichum coccodes</i>)	Awaiting results									
Black scurf (<i>Rhizoctonia solani</i>)	Awaiting results									
Dry rot (<i>Fusarium coeruleum</i>)									6	
Dry rot (<i>Fusarium sulphureum</i>)										9
Internal damage (bruising)					5					
External damage (and splitting)									6	

FIELD AND STORE QUALITIES	LOW	→								HIGH
Drought resistance									7	
Heat tolerance									6	
Storability									7	

Disclaimer: The contents are based on the best information available, but the selling parties will not accept responsibility for any inaccuracy caused by circumstances outside their control, nor will they be liable for any costs, loss, damage or injury resulting from interpretation of, or decisions based on, the information provided.